

Coastal AgroBusiness, Inc.



Product Label



Liquid Carbon
Soil Conditioner

| | |
|------------------------|------|
| Organic Carbon | 5% |
| Inert Ingredients..... | 95% |
| ----- | |
| Total | 100% |

Derived from completely organic Leonardite.
Non-Plantfood Ingredients

CAUTION

KEEP OUT OF REACH OF CHILDREN

Read entire label for additional precautionary statements.

SHAKE WELL BEFORE USING

Manufactured By:
Coastal AgroBusiness, Inc
Greenville, North Carolina

CarboMaxx is a highly concentrated product designed for direct field application, injection into irrigation systems, or for addition to liquid fertilizers used as starter solutions. CarboMaxx contains materials that aid in the formation of soil particles and play a vital role in helping to build organic matter in soils. CarboMaxx contains natural humic acids and improves nutrition and yields in a wide variety of crops. CarboMaxx helps relieve plant stress whether it's applied as a soil amendment or mixed with fertilizer. Natural bio-stimulants found in CarboMaxx improve the availability of nutrients in the soil, particularly phosphorus and micronutrients. The soils nutrient holding capacity is increased as well as water holding capacity. The workability of problem soils is improved over time and weak areas in a field become less noticeable. CarboMaxx enhances any crops ability to absorb nutrients and resist stress, especially in marginal soils, resulting in faster growth and higher yields.

Note: Be careful when mixing CarboMaxx with other products. Check product compatibilities in small amounts before large scale mixing. Do not mix CarboMaxx with ammonium sulfate nitrogen sources.

PRECAUTIONARY STATEMENTS / HAZARDS TO HUMANS

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Can cause skin, mucous membrane and eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mists. Use only with adequate ventilation. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

Do not apply this product in a manner as to directly or through drift expose workers or other persons. Always read and follow product labeling.

STATEMENT OF PRACTICAL TREATMENT

If Inhaled: Remove to fresh air. Give artificial respiration if necessary. If breathing is difficult, get immediate medical attention.

If Swallowed: If conscious, give large amounts of water. Do not induce vomiting. Get immediate medical attention. Never induce vomiting or give anything by mouth to an unconscious person.

If In Eyes: Flush eyes with copious amounts of clean water for at least 15 minutes. Get medical attention.

If On Skin: Remove contaminated clothing and wash skin with soap and water. If irritation develops, get medical attention.

NOTE: If contact is made with spray solution containing pesticides, follow the "First Aid" statement on the pesticide label.

STORAGE AND DISPOSAL

Do not store near heat or open flame. Store in original container only. Keep container tightly closed and do not allow water to be introduced into the contents of this container. Keep from freezing. If damaged or leaking, soak up with absorbent material and place in suitable container for disposal. Transfer remaining product to suitable container for storage and label.

Do not contaminate water, food or feed by storage or disposal. Do not contaminate water by runoff from equipment wash waters or spray materials. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Triple rinse (or equivalent) and add rinsate to spray tank, then offer for recycling, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

AGRICULTURAL USE REQUIREMENTS AND USER SAFETY RECOMMENDATIONS

When used in conjunction with pesticides, consult pesticide manufacturer's label for worker protection recommendations. Use of waterproof gloves and protective eyewear is recommended when handling this product.

DIRECTIONS FOR USE

General use is to add CarboMaxx to liquid fertilizers at the rate of 1-2 percent of the final solution. Use a total rate for the growing season of 2-10 quarts per acre. An initial application of 2 quarts per acre should be applied early in the season before or after a major fertilizer application. The remainder should be split into several applications made during the growing season. The higher rates should be used for specific problem areas, such as nematode infestations or soil reclamation.

Note: Be careful when mixing CarboMaxx with other products. Check product compatibilities in small amounts before large scale mixing. Do not mix CarboMaxx with ammonium sulfate nitrogen sources.

RATE SELECTION CHART

| Soil Application (All Crops) | RATE |
|-------------------------------|---------------------------------|
| Direct Application to Soil | 2 - 4 quarts/acre |
| Addition to Starter Solutions | 1 – 2 percent of final solution |
| Injection or Drip Irrigation | 2 – 4 quarts per 100 gallons |

SOIL APPLICATION

Direct Application to Soil: CarboMaxx may be broadcast directly onto soil. Apply in a minimum of 15 gallons of water per acre. For best results, CarboMaxx should be incorporated. Do not apply in solutions containing sulfur.

Addition to Starter Solutions: CarboMaxx should be added to starter at 1% – 2% of final solution (1-2 gallons per 100 gallons of solution). May be used in furrow, banded 2x2, or on top of row in addition to normal at-plant application.

Injection or Drip Irrigation: CarboMaxx should be added at a rate of 2-4 quarts per 100 gallons. Addition should be made several times during the growing season. Can be used as an additive to your normal drip fertigation schedule.

CarboMaxx provides a soluble source of humus that improves the organic matter fraction of the soil. Crop production is increased as the efficiency of N-P-K fertilizers is improved and micronutrients found in the soil are activated for plant use. CarboMaxx is an excellent additive for starter fertilizers, particularly when micronutrients such as zinc, manganese and iron are being applied. Tests show that plants grow faster and more robustly where CarboMaxx is used on sodic or calcareous soils or on soils low in organic matter.

For pot drenching

Use a one percent solution and enough solution to go through the soil (dripping at the bottom).

For an established lawn:

Apply 8 ounces per 1000 square feet as growth begins in the spring and then repeat the application early in the summer to reduce compaction and improve soil tilth and water holding capacity.

- Use a hose proportioner to apply, using plenty of water to incorporate the material into the soil.
- Or spray it on and irrigate it in with 1/2 inch of water.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions contained on this label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to damage to plants, crops, and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.